

NICE monitoring guidance for DOACs

| | | Repeat every: | | | Renal and liver function tests should be performed more often if there is an intercurrent illness that may impact renal or hepatic function. |
|-------------|----------------------------|----------------------------------|---|-----------|--|
| DOACs | At the start of treatment: | 3 months | 6 months | 12 months | |
| Apixaban | Baseline clotting screen | | | | |
| | renal function tests | ✓ if (CrCl) between 15-30 mL/min | ✓ if (CrCl) between 30-60 mL/min | ✓ | |
| | liver function tests | | | ✓ | |
| | a full blood count | | | ✓ | |
| Rivaroxaban | Baseline clotting screen | | | | |
| | renal function tests | ✓ if (CrCl) between 15-30 mL/min | ✓ if (CrCl) between 30-60 mL/min | ✓ | |
| | liver function tests | | | ✓ | |
| | a full blood count | | | ✓ | |
| Edoxaban | Baseline clotting screen | | | | |
| | renal function tests | | ✓ if (CrCl) between 30-60 mL/min, is above 75years old, and/or is fragile | ✓ | |
| | liver function tests | | | ✓ | |
| | a full blood count | | | ✓ | |
| | Baseline clotting screen | | | | |

| | | | | |
|-------------------|----------------------|--|---|---|
| Dabigatran | renal function tests | | ✓ if (CrCl) between 30-60 mL/min, is above 75years old, and/or is fragile | ✓ |
| | liver function tests | | | ✓ |
| | a full blood count | | | ✓ |

All information from the following link:

<https://cks.nice.org.uk/anticoagulation-oral>